



Research Article

Chemical composition of the essential oil of *Vernonia gracilis* Kunth from Venezuela

Alida Pérez-Colmenares* , Luis Rojas-Fermín , Rosa Aparicio-Zambrano  and Ysbelia Obregón-Díaz 

Research Institute "Dr. Alfredo Nicolás Usubillaga del Hierro", Faculty of Pharmacy and Bioanalysis, University of Los Andes, Mérida, C.P. 5101, Venezuela.

Article Information

Received: 20 December, 2023
Revised: 10 February, 2023
Accepted: 16 February, 2023
Published: 23 February, 2023

Academic Editor

Prof. Dr. Radosław Kowalski

Corresponding Author

Alida Pérez-Colmenares
E-mail: alidaperez@gmail.com
Tel.: +58 4167741507

Keywords

Vernonia; *Vernonia gracilis*,
essential oil, β -caryophyllene,
germacrene D, bicyclogermacrene.

Abstract

The essential oil from the *Vernonia gracilis* Kunth (Asteraceae) leaves collected in San Juan de Lagunillas, Venezuela was obtained by hydrodistillation and its composition was determined by GC/MS. The yield of the essential oil was 0.001 % (v/w), calculated on a dry weight basis. Twenty four compounds were identified, representing 93.19 % of the oil. The most abundant components were β -caryophyllene (23.15 %), germacrene-D (18.68 %), bicyclogermacrene (18.31 %), α -humulene (6.38 %), caryophyllene epoxide (4.84 %) and spathulenol (4.53 %).

1. Introduction

Vernonia (family Asteraceae, tribe Vernonieae) is a genus of about 1300 species, it's widely distributed in tropical and sub-tropical areas, especially in South America, Asia and Africa [1]. Several *Vernonia* species have been used in traditional medicine to treat reproductive problems, including urinary tract infections, diabetes mellitus, bacterial infections and malaria [2-6]. Various studies have reported antioxidant, anthelmintic, antimicrobial, anti-diabetes, anti-inflammatory, antitumor, analgesic, anti-plasmodial, anticancer and neuroprotective activities [7-11].

Numerous phytochemical studies of the genus *Vernonia* have revealed the presence of diterpenes, triterpenes, steroids, and the most frequently isolated chemical constituents are flavonoids and sesquiter-

pene lactones, which have been used as taxonomic markers [12-16]. Despite this extensive investigation, little is known about the composition of the genus *Vernonia* essential oils.

Vernonia gracilis Kunth is a synonym for *Lepidaploa gracilis* [17,18]. Literature surveys reveal that no phytochemical investigations have been previously conducted on the essential oil of *V. gracilis*, the present study was undertaken to obtain knowledge of their chemical composition.

2. Materials and methods

2.1 Plant material

Leaves of *V. gracilis* were collected in May 2016 at San Juan de Lagunillas, Mérida State, Venezuela. A voucher specimen (LCGM01) has been deposited at

Table 1. Percentage composition of the essential oil from leaves of *Vernonia gracilis* Kunth.

No.	Compounds	Composition of essential oil (%)	^a RI	^b RI
1	β -Pinene	0.46	965	979
2	Limonene	0.16	1032	1029
3	<i>Cis</i> -3-hexenyl 3-methylbutanoato	0.19	1231	1235
4	β -Copaene	2.18	1382	1387
5	β -Bourbonene	0.74	1385	1388
6	β -Elemene	0.98	1396	1390
7	β -Gurjunene	0.35	1428	1433
8	β-Caryophyllene	23.15	1440	1433
9	Epi-Bicyclosesquiphellandrene	0.34	1448	1463
10	α-Humulene	6.38	1457	1465
11	Alloaromadendrene	2.10	1471	1460
12	α -Amorphene	0.75	1482	1484
13	Germacrene-D	18.68	1492	1485
14	Bicyclogermacrene	18.31	1502	1500
15	Germacreno A	0.77	1503	1509
16	γ -Cadinene	0.60	1513	1528
17	δ -Cadineno	1.56	1526	1530
18	Spathulenol	4.53	1575	1578
19	Caryophyllene epoxide	4.84	1603	1588
20	Viridiflorol	1.45	1611	1610
21	Humulene epoxide	0.63	1609	1608
22	Isospathulenol	1.27	1624	1626
23	α -Cadinol	1.68	1661	1672
24	T-Muurolol	1.09	1658	1659
Total Identified		93.19		

^aRI: Retention index this work. ^bRI: Theoretical Retention index.

the MERF Herbarium (Herbarium of the Faculty of Pharmacy and Bioanalysis in Mérida, Venezuela). The botanical identification was made by Dr. Pablo Meléndez.

2.2 Extraction and analysis of the essential oil

Fresh leaves (1000 g) were cut into small pieces and submitted to hydrodistillation for 3 hours using a Clevenger-type apparatus. A volume of 0.1 mL of essential oil was obtained (0.001 %, v/w). The composition of the essential oil was determined by comparison of the mass spectrum of each component with Wiley GC/MS library data and also from RI data.

2.3 Gas chromatography – mass spectrometry

Gas chromatography – mass spectrometry (GC-MS) analyses were conducted on a Model 5973 Hewlett-Packard GC-MS system fitted with a HP- 5MS fused silica column (30 m x 0.25 mm i.d., film thickness 0.25 μ m, Hewlett-Packard). The oven temperature program was the same as the one used for the HP-5 column for GC analysis; the transfer line temperature was programmed from 150 °C to 180 °C; source

temperature, 230 °C; quadrupole temperature 150 °C; carrier gas, helium adjusted to a linear velocity of 34 cm/s; ionization energy, 70 eV; scan range, 40 to 500 amu; 3.9 scans/s. A sample of a 2 % solution of the essential oil (1.0 μ L) was injected using a Hewlett-Packard ALS injector with a split ratio of 50:1. The identity of the oil components was established from their GC retention indices, by comparing their MS and those of standard components available in the laboratory, and through library search (Nist 05 and Wiley MS Data Library, 6th edn) [19, 20].

3. Results and discussion

The hydrodistillation from the leaves of *V. gracilis* produced 0.1 mL yellow oil with a yield of 0.001 %. The chemical composition of the oil was investigated using GC-MS technique. Twenty-four components were identified in the leaves' oil, which represent 93.19 % of the total oil. These compounds and their retention indices (RI) and relative percentage concentrations are listed in Table 1 and Fig. 1,




Figure 1. GC chromatogram (HP-5 column) and representative compounds of leaves of *Vernonia gracilis* Kunth.

according to the elution order on HP-5 column. The identified products may be divided into four different groups: monoterpenes (0.62 %), oxygenated monoterpenes (0.19 %), sesquiterpenes (76.89 %) and oxygenated sesquiterpenes (15.49 %). The main components of the leaves' oil were β -caryophyllene (23.15 %), germacrene-D (18.68 %), bicyclgermacrene (18.31 %), α -humulene (6.38 %), caryophyllene epoxide (4.84 %) and spathulenol (4.53 %). This is the first report on the volatile compounds of *V. gracilis*, the present data could not be compared with other analysis of the same species.

There are several reports on the oil composition of the plants of the genus *Vernonia* described in other parts of the world (Vietnam, Brazil, India and Nigeria) [21]. Most of them are characterized by the presence of monoterpenoids and sesquiterpenoids. The preliminary chemical analysis of the essential oils of *Vernonia* has been performed earlier the major components were β -caryophyllene (28.50 %), caryophyllene oxide (16.60 %), α -copaene (9.00 %) and α -humulene (7.10 %) from the leaves of *V. patula* [21] and the essential oil was obtained from *V. chalybaea* showing it was rich in β -caryophyllene (39.06 %) and bicyclgermacrene (19.69 %) [22]. The major components found in the essential oil from aerial parts of *V. scorpiodes* were β -caryophyllene (30.60 %), germacrene D (27.30 %), and bicyclgermacrene (8.50 %) [23]. On the other hand, the major components of the essential oils from the aerial parts of *V. amygdalina* and from the roots of *V. cinerea* were α -muurolol (45.70 %) and α -muurolene (30.70 %),

respectively [24, 25]. The principal components of the essential oil analysis of *V. crotonoides* were α -cadinol (14.10 %), spathulenol (9.80 %), caryophyllene oxide (5.90 %), and epi- α -cadinol (5.30 %) [26].

The major constituents reported in the essential oils from different organs of *V. condensata* were the leaf oil β -caryophyllene (18.35 %), γ -muurolene (16.41 %) and α -humulene (6.03). Linalol (12.13 %), 10-epi-italicene ether (8.88 %), myrcene (7.68 %), and 2- δ -carene (6.86 %) were in the stem oil. The monoterpenes 1,8-cineole (11.03 %), thymol (7.68 %) and linalool (6.30 %) were the major constituents reported in the flower oil [27]. It is well known that sesquiterpenes compounds, as identified in the present study, predominate in the essential oils of the genus *Vernonia*. These differences are mainly due to variations in the ecological conditions, genetic and environmental factors, such as temperature, moisture, the soil, the harvest period, geographical position, seasonal variations and the vegetative cycle are important [28, 29].

4. Conclusions

In the current investigation, the chemical composition of essential oil of fresh leaves of *Vernonia gracilis* (Asteraceae) collected in Mérida, Venezuela was evaluated. The essential oil is dominated by the presence of sesquiterpenes (76.89 %) and oxygenated sesquiterpenes (15.49 %). The main components were β -caryophyllene (23.15 %), germacrene-D (18.68 %) and bicyclgermacrene (18.31 %). To our knowledge, this is the first time that the chemical composition of the essential oil of the leaves *V. gracilis* of has been

reported and these compounds may also aid in future chemotaxonomic research of this species.

Authors' contributions

Conceptualization, L.R.F.; Methodology, L.R.F.; Software, L.R.F.; Validation, A.P.C., Formal analysis, A.P.C.; Y.O.D.; Investigation, A.P.C., R.A.Z., Y.O.D.; Resources, A.P.C., R.A.Z., Y.O.D.; Data curation, R.A.Z.; Writing – original draft preparation, A.P.C., Y.O.D.; Writing – review & editing, A.P.C., R.A.Z., Y.O.D.

Acknowledgements

All authors are grateful to the Research Institute “Dr. Alfredo Nicolás Usabillaga del Hierro”, Faculty of Pharmacy and Bioanalysis, University of Los Andes, Mérida 5101, Venezuela.

Funding

This research received no specific grant from any funding agency.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Toyang, N.; Verpoorte, R. A review of the medicinal potentials of plants of the genus *Vernonia* (Asteraceae). *J Ethnopharmacol.* 2013, 146, 681-723. <https://doi.org/10.1016/j.jep.2013.01.040>.
2. Focho, D.A.; Nkeng, E.A.P.; Lucha, C.F.; Ndam, W.T.; Afegenui, A. Ethnobotanical survey of plants used to treat diseases of the reproductive system and preliminary phytochemical screening of some species of Malvaceae in Ndop Central Sub-Division, Cameroon. *J. Med. Plants Res.* 2009, 3, 301–314.
3. Onyibe, P.N.; Edo, G. I.; Nwosu, L. C.; Ozgor, E. Effects of *Vernonia amygdalina* fractionate on glutathione reductase and glutathione-S-transferase on alloxan induced diabetes wistar rat. *Biocatal. Agric. Biotechnol.* 2021, 36, 102-118. <https://doi.org/10.1016/j.bcab.2021.102118>.
4. Jepkoech, C.; Omosa, L.K.; Nchiozem-Ngnitedem, V.A.; Kenanda, E.O.; Guefack, M.G.; Mbaveng, A.T.; Heydenreich, M. Antibacterial secondary metabolites from *Vernonia auriculifera* Hiern (Asteraceae) against MDR phenotypes. *Nat. Prod. Res.* 2022, 36, 3203-3206. <https://doi.org/10.1080/14786419.2021.1953024>.
5. Collins, S.; Mouthé, G.; Nguiam, M.; Tchamgoue, J.; Jouda, J.B.; Longo, F.; Fogue, S. Antibacterial flavonoids and other compounds from the aerial parts of *Vernonia guineensis* Benth. (Asteraceae). *Chem. Biodivers.* 2020, 17, e2000296. <https://doi.org/10.1002/cbdv.202000296>.
6. Bihonegn, T.; Fentahun, S. Review on malaria and antimalarial activity of *Vernonia amygdalina* in Ethiopia: A review article. *J. Nurs. Healthc. Res.* 2019, 60, 1-15. <https://doi.org/10.7176/jhmn/60-01>.
7. Osei, C.; Acheampong, A.; Boakye, Y. D.; Asante, B.; Ohene, S.; Amankwah, F. Anthelmintic, anti-inflammatory, antioxidant, and antimicrobial activities and FTIR analyses of *Vernonia camporum* stem-bark. *J. Chem.* 2021, 2021. <https://doi.org/10.1155/2021/3328073>.
8. Asante, D.B.; Henneh, I.T.; Acheampong, D.O.; Kyei, F.; Adokoh, C. K.; Ofori, E. G.; Ameyaw, E.O. Anti-inflammatory, anti-nociceptive and antipyretic activity of young and old leaves of *Vernonia amygdalina*. *Biomed. Pharmacother.* 2019, 111, 1187-1203. <https://doi.org/10.1016/j.biopha.2018.12.147>.
9. Zemicheal, G.; Yalemtehay, M. Antiplasmodial activity of *Vernonia adoensis* aqueous, methanol and chloroform leaf extracts against chloroquine sensitive strain of *Plasmodium berghei* *in vivo* in mice. *BMC Res. Notes.* 2018, 11, 1-6. <https://doi.org/10.1186/s13104-018-3835-2>.
10. Hasibuan, P.; Harahap, U.; Sitorus, P.; Satria, D. The anticancer activities of *Vernonia amygdalina* Delile. Leaves on 4T1 breast cancer cells through phosphoinositide 3-kinase (PI3K) pathway. *Heliyon.* 2020, 6, e04449. <https://doi.org/10.1016/j.heliyon.2020.e04449>.
11. Oboh, G.; Adedayo, B.C.; Adetola, M.B.; Oyeleye, I.S.; Ogunsuyi, O.B. Characterization and neuroprotective properties of alkaloid extract of *Vernonia amygdalina* Delile in experimental models of Alzheimer's disease. *Drug Chem. Toxicol.* 2022, 45, 731-740. <https://doi.org/10.1080/01480545.2020.1773845>.
12. KTurak, A.; Maimaiti, Z.; Aisa, H. A new kaurane diterpene from the seeds of *Vernonia anthelmintica*. *Chem. Nat. Compd.* 2018, 54, 475-477. <https://doi.org/10.1007/s10600-018-2382-3>.
13. Zemene, T.; Meshesha, M.; Denu, D.; Abdissa, N. Tetracyclic triterpenes from the stem bark of *Vernonia bialfrae* and evaluation of its antimicrobial activity. *Ethiop. J. Agric. Sci.* 2020, 15, 56-64.
14. Liu, X.; Tian, W.; Wang, G.; Xu, Q.; Zhou, M.; Gao, S.; Chen, H. Stigmastane-type steroids with unique conjugated $\Delta^7, 9$ (11) diene and highly oxygenated side chains from the twigs of *Vernonia amygdalina*. *Phytochemistry.* 2019, 158, 67-76. <https://doi.org/10.1016/j.phytochem.2018.10.036>.
15. Alara, O.R.; Abdurahman, N.H.; Olalere, O.A. Ethanolic extraction of flavonoids, phenolics and antioxidants from *Vernonia amygdalina* leaf using two-level factorial design. *J. King Saud Univ. Sci.* 2020, 32, 7-16. <https://doi.org/10.1016/j.jksus.2017.08.001>.

16. Bitchagno, G.T.; Schuffler, A.; Gross, J.; Krumb, M.; Tane, P.; Opatz, T. Sesquiterpene lactones from *Vernonia tufnelliae*: Structural characterization and biological evaluation. *J. Nat. Prod.* 2022, 85, 1681-1690. <https://doi.org/10.1021/acs.jnatprod.2c00055>.
17. Ávila, F.; Funk, V.A.; Diazgranados, M.; Díaz-Piedrahíta, S.; Vargas, O. Asteraceae. En: R. Bernal, S. R. Gradstein y M. Celis (Eds.), *Catálogo de plantas y líquenes de Colombia*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá. 2015.
18. Pruski, J.F. Compositae of Central America–VIII. The genus *Lepidaploa* (Vernonieae). *Phytoneuron*. 2017, 50, 1–39.
19. Adams, R.P. Identification of essential oil components by gas chromatography/mass spectroscopy. 4th Edn.; Allured Publishing Corp., Carol Stream, Illinois. 2017.
20. Davies NW. Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and carbowax 20 M. phases. *J. Chromatogr. A*. 1990, 503, 1-24. [https://doi.org/10.1016/s0021-9673\(01\)81487-4](https://doi.org/10.1016/s0021-9673(01)81487-4).
21. Hoi, T.M.; Chung, N.T.; Huong, L.T.; Ogunwande, I.A. Studies on Asteraceae: Chemical compositions of essential oils and antimicrobial activity of the leaves of *Vernonia patula* (Dryand.) Merr. and *Grangea maderaspatana* (L.) Poir. from Vietnam. *J. Essent. Oil-Bear. Plants*. 2021, 24 500-509. <https://doi.org/10.1080/0972060x.2021.1947390>.
22. Nogueira, A.C.; Morais, S.M.; Souza, E.B.; Albuquerque, M.R.; Santos, H.S.; Cavalcante, C.S.; Fontenelle, R.O. Antifungal and antioxidant activities of *Vernonia chalybaea* Mart. ex DC. essential oil and their major constituent β -caryophyllene. *Braz Arch Biol Technol*. 2020, 63. <https://doi.org/10.1590/1678-4324-2020190177>.
23. Albuquerque, M.R.; Lemos, T.L.; Pessoa, O.D.; Nunes, E.P.; Nascimento, R.F.; Silveira, E.R. Chemical composition of the essential oil from *Vernonia scorpioides* (Asteraceae). *Flavour Fragr. J.* 2007, 22 249-250. <https://doi.org/10.1002/ffj.1782>.
24. Ogunbinu, A.O.; Flamini, G.; Cioni, P.L.; Ogunwande, I.A.; Okeniyi, S.O. Essential oil constituents of *Eclipta prostrata* (L.) L. and *Vernonia amygdalina* Delile. *Nat. Prod. Commun.* 2009, 4 1934578X0900400321. <https://doi.org/10.1177/1934578x0900400321>.
25. Joshi, R.K. GC/MS analysis of the essential oil of *Vernonia cinerea*. *Nat. Prod. Commun.* 2015, 10, 934578X1501000746. <https://doi.org/10.1177/1934578x1501000746>.
26. De Carvalho, C.C.; Turatti, I.C.; Lopes, N.P.; Do Nascimento, A.M. Chemical composition of the essential oil of *Vernonia crotonoides*. *Chem. Nat. Compd.* 2013, 49, 761-762. <https://doi.org/10.1007/s10600-013-0734-6>.
27. Lopes, C.B.; Da Camara, C.A.; De Moraes, M.M. Composition of essential oils from the leaves, stems, and flowers of *Vernonia condensata* of Pernambuco, Brazil. *Chem. Nat. Compd.* 2019, 55, 756-758. <https://doi.org/10.1007/s10600-019-02802-8>.
28. Gobbo, N.; Lopes, N. Plantas medicinais: Fatores de influência no conteúdo de metabólitos secundários. *Quim. Nova*. 2007, 30, 274-381. <https://doi.org/10.1590/s0100-40422007000200026>.
29. Figueiredo, A.; Barroso, L.; Scheffer, P. Factors affecting secondary metabolite production in plants: volatile components and essential oils. *Flavour Fragr. J.* 2008, 23, 213-226. <https://doi.org/10.1002/ffj.1875>.