











Research Article

Piper robustipedunculum Yunck.: Essential oil profile and chemosystematic insight of this Brazilian native species

Ygor Jessé Ramos^{1,2*} , Irene Candido Fonseca¹ , George Azevedo Queiroz¹ , Daniel de Brito Machado^{1,2} , Claudete da Costa-Oliveira¹ , Jéssica Regina Sales Felisberto¹ , Elsie Franklin Guimarães¹  and Davyson de Lima Moreira^{1,2*} 

1. Botanical Garden Research Institute of Rio de Janeiro, Jardim Botânico, Rio de Janeiro, RJ, 22460-030, Brazil.
2. Rio de Janeiro State University, Rio de Janeiro, RJ, Maracanã, 20550-013, Brazil.

Article Information

Received: 24 January, 2023
Revised: 13 February, 2023
Accepted: 17 February, 2023
Published: 24 February, 2023

Academic Editor

Prof. Dr. Radosław Kowalski

Corresponding Authors

Prof. Dr. Ygor Jessé Ramos
ygorjesse@jbrj.gov.br
Prof. Dr. Davyson de Lima Moreira
davysonmoreira@jbrj.gov.br

Keywords

Aromatic plant, chemophenetic, Piperaceae, *E*-nerolidol, hydrodistillation, sesquiterpenes.

Abstract

The aim of this study was to analyze the chemical composition of the essential oil extracted from the leaves of the species *Piper robustipedunculum* Yunck. The essential oil was obtained through hydrodistillation and analyzed using GC-FID and GC-MS. The results showed that 93.88% of the total composition was composed of 24 compounds. The essential oil was light yellow in color, with a yield of 0.68%, and was rich in sesquiterpenes, including both oxygenated (37.09%), such as *E*-nerolidol (13.60%), and non-oxygenated sesquiterpenes (45.06%), such as δ -cadinene (10.72%), α -copaene (9.85%), and valencene (8.34%) as the major components. This essential oil has the potential for industrial and pharmaceutical applications, and can aid in future research in ecology, chemotaxonomy, and chemophenetics related to the Piperaceae species.

1. Introduction

Piperaceae family belongs to the Piperales order, which is part of the Magnoliidae group. This pantropical family encompasses approximately 4,100 plant species found worldwide, with the largest genus being *Piper* L., containing around 2,000 species [1]. In Brazil, the highest diversity of this genus is located in the Atlantic and Amazon Forests, housing 347 taxa out of the 466 present in Brazil [1]. The Northeast region of Brazil records 93 species of this genus.

Several well-known plants from the *Piper* genus, such as Black Pepper (*Piper nigrum* L.), Long Pepper (*Piper longum* L.), and Kava-Kava (*Piper methysticum* G.

Forst), are commonly used as condiments and have various therapeutic purposes [4, 5]. Ethnobotanical surveys of medicinal plants emphasize the importance of the *Piper* genus for its medicinal properties, including increased gastrointestinal motility, antifungal, antibacterial, larvicidal, antinociceptive, and antihepatotoxic activities. The popularity of its medicinal use is linked to the number of *Piper* species available [6-7].

Due to their similar morphological features, species from this genus can often be found in the presence of different ethnospecies, including Jaborandi (*Piper*

amalago L., *Piper crassinervium* HB & K., *Piper mollicomum* Kunth and *Piper aduncum* L.), Pariparoba (*Piper cernuum* Vell., *Piper mikianium* (Kunth) Steudel and *Piper umbellatum* L.), White Betis (*Piper rivinoides* Kunth., *Piper arboreum* Aubl. and *Piper tuberculatum* Jacq.), among others [8-9].

The essential oils extracted from various parts of the *Piper* genus have demonstrated remarkable biological properties, including antioxidant, antibacterial, antifungal, antiprotozoal, antiproliferative, anti-inflammatory, and neuropharmacological activities [10-12, 7, 13]. This diversity of activities can be attributed to the extensive chemical diversity present in *Piper*'s essential oils, which make them valuable tools for chemophenetic and ecological characterization, particularly for new and unstudied species. The presence of specialized secondary metabolites in these oils provides important information for taxonomic descriptions and suit for defining characteristic of a particular species [14].

Piper robustipedunculum Yunck (Fig. 1) is a species native to Brazil, restricted to the Northeastern states. It has morphological similarities with *Piper amplum* Kunth, *Piper arboreum* Aubl. and *Piper ilheusense* Yunck which are plants of popular use and coexist in the same region [15-18]. However, this species has no phytochemical approach in the literature.



Figure 1. Habit of *Piper robustipedunculum* Yunck.

With all the above, this work aims to characterize the composition of the essential oil of *Piper robustipedunculum* Yunck from a natural population of the city of São Vicente Férrer, Pernambuco, Northeastern Brazil.

2. Materials and methods

2.1 Plant material and isolation of essential oil

Piper robustipedunculum Yunck leaves were collected (three samples) on October 22, 2019 at 9 am, in the municipality of São Vicente Férrer, at coordinates 7° 36'51" N and 35° 30' 53" E, at an elevation of 352 m.d. An exsiccate was identified and deposited in the Herbarium of the Botany Department of the National Museum of the Federal University of Rio de Janeiro under voucher number 45321. This study was assigned in the Brazilian Genetic Heritage Management Council under the number AE4E953.

The hydrodistillation was the chosen extraction technique, with Clevenger apparatus type, where 100 g of fresh leaves was placed in a volumetric flask (2 L) with 700 mL of distilled water and heated for 2 h. Yielding was calculated in relation to the obtained essential oil (g) and the amount of fresh leaves (g). The essential oil was kept in sealed amber flasks under refrigeration -20 °C until analysis [12-13].

2.2 Essential oil analysis

The chemical characterization and quantification of *P. robustipedunculum* essential oil was made by gas chromatography coupled to mass spectrometry (GC-MS) and a gas chromatography equipped with a flame ionization detector (FID), respectively. Samples in triplicates were diluted in dichloromethane (HPLC grade, Tedia, Brazil) before analysis (1 mg/mL).

A 1 µL of the solution was injected into an HP Agilent GC 6890 coupled to Agilent MS 5973 series mass selective detectors, splitless mode, wherein the injector temperatures were set at 270°C, operating at 70 eV in positive mode. A HP-5MS (5%-phenylmethylpolysiloxane) capillary column [Agilent J & W; GC Columns (USA)] was used with 30 m x 0.25 mm i.d. x 0.25 µm particle size. Chromatography conditions were started at 60 - 240 °C at 3 °C / min, totaling 60 min. For the separation of the constituents, helium (~99.9999%) was used as a carrier gas at 1,0 mL/ min, at a rate of 1,0 s sweeps and mass range of *m/z* 40 - 600 atomic mass unit (*u*) [12-13].

The GC-FID was achieved in a chromatograph equipped with a flame ionization detector [HP-Agilent 6890 GC-FID]. A 1 µL of the solution were injected under the same analytical conditions described above, except for the carrier gas used,

which was hydrogen at flow rate of 1,0 mL/min. The retention times (Rt) of the compounds were measured in minutes and they were used to calculate their linear retention index, obtained from the injection of a homologous series of hydrocarbons (C₈-C₂₅ de *n*-alkane, Sigma-Aldrich, Brazil) under the same sample analytical condition [19-20, 12-13].

The mass spectra of the constituents were compared with those from library (NIST, 98 e WILEY 7n) and with those published in the literature [21]. Additionally, when possible, the samples were analyzed in conjunction with authentic standards (Sigma-Aldrich, St. Louis, MS, USA).

3. Results and discussion

The essential oil (EO) obtained from the leaves of *P. robustipedunculum*, a native species and endemic to Northeastern Brazil, collected in the region of the Atlantic Forest and studied by GC-MS and GC-FID are presented in Table 1. A chromatographic profile of the essential oil is also presented in Fig. 2. EOs obtained by the same methods in other *Piper* species also showed high percentage of monoterpenes and sesquiterpenes in leaves and reproductive parts in all investigation periods [14].

The EO showed light-yellow color and average yield of 0.68% (g/g). This is very important to industrial applications when compared to other commercial species [14, 22-23].

It was possible to characterize 93.88% of the total EO. Sesquiterpenes (acyclic, monocyclic, bicyclic and tricyclic) were registered as the main fraction, with a total of 82.15%, being non-oxygenated 45.09% and oxygenates 36.09%. *E*-nerolidol (13.60%), δ -cadinene (10.72%), α -copaene (9.85%), and valencene (8.34%) were identified as major compounds. For Piperaceae species, the high content of sesquiterpenes is in accordance with literature data [24, 7]. The monoterpene class represent only 2.66% of the total. These similar variations of monoterpenes and sesquiterpenes have already been reported by Santos et al. [24] in Brazilian Atlantic Forest for species of the same genus, including *Piper arboreum* var. *arboreum* Yunck. and *Piper goesii* Yunck. The Piperaceae family is known for the frequent occurrence of ethnospices, as previously mentioned.

Considering taxonomic approaches, in the subclade Isophyllon [25], the taxonomic classification group that belongs to *P. robustipedunculum*, it is possible to

Table 1. Aromatic profile of *P. robustipedunculum* essential oil from leaves (*n* = 3) from Pernambuco, Brazil.

RT	Compounds ¹	RI*	RI**	Composition (%)
9.792	limonene	1026	1024	1.05
13.440	Z- β -ocimene	1034	1032	1.61
22.002	δ -elemene	1323	1335	1.62
24.043	α -copaene	1374	1374	9.85
25.679	<i>E</i> -caryophyllene	1419	1417	4.75
26.916	α -humulene	1455	1452	1.05
27.800	<i>E</i> -cadin-1(6),4-diene	1476	1475	1.58
27.972	germacrene D	1483	1480	2.43
28.302	β -selinene	1490	1489	1.12
28.746	valencene	1500	1496	8.34
28.816	γ -cadinene	1515	1513	4.60
29.911	δ -cadinene	1524	1522	10.72
31.163	<i>E</i> -nerolidol	1560	1561	13.60
32.163	germacrene D-4-ol	1574	1574	5.16
32.359	glenol	1587	1586	1.02
32.649	viridiflorol	1593	1592	1.54
33.229	1- <i>epi</i> -cubenol	1628	1627	1.94
33.362	Z-cadin-4-en-7-ol	1634	1635	1.04
33.752	<i>epi</i> - α -cadinol	1640	1638	2.21
33.960	<i>epi</i> - α -muurolol	1642	1640	2.17
35.010	α -muurolol	1645	1644	2.12
34.519	α -cadinol	1657	1652	5.29
35.796	tricosane	2299	2300	3.98
48.573	octacosane	2798	2800	4.09
Total identified				93.88
Compound classes (%)				
Non-oxygenated Monoterpene				2.66
Non-oxygenated Sesquiterpene				45.06
Oxygenated sesquiterpenes				37.09
Others (hydrocarbons)				8.07

¹Compounds an in order of elution on the HP-5MS column; RT = Retention time in minutes; RI* = Experimentally determined retention indices for C₈ - C₂₀ *n*-alkanes on the HP-5MS RI column; RI** = Literature Retention Indices [21], %= average compound concentration in the three samples obtained.

find species that in their EO from leaves compounds produced by acetate-mevalonate or methylerythriol-4-phosphate metabolic pathways.

In 1966, Yuncker [26] pointed out the ease of errors caused by the similarity in morphology among the species within the Isophyllon. Specifically, the




Figure 2. Chromatographic profile of the flame ionization detector (CG-FID) of the essential oil of *P. robustipedunculum* from Pernambuco, Brazil. (See compound 1 - α -copaene; 2 – valencene; 3 - δ -cadinene; and 4 - *E*-nerolidol)

medicinal plants *Piper amplum* Kunth, *Piper arboretum* Aubl., *Piper ilheusense* Yunck and *Piper vicosanum* Yunck are often confused with *P. robustipedunculum* due to their similar appearance, which can lead to accidental or intentional mislabeling in Northeast Brazil. This can be a problem for quality control and the correct use of these plants [9, 18].

Chemotaxonomically allow for the observation of variations in chemical compositions among species within this group across different studies in the literature. For example, for the EO of *Piper ilheusense* Yunck it was registered the compounds patchouli alcohol (11.10%), *E*-caryophyllene (11.80%), gleenol (7.50%) and δ -cadinene (6.90%) [16]. Pereira et al. [27] describe the chemical composition of EO from different species of *Piper*, in which, *Piper amplum* contained the iso-leptospermane (26.44%) and α -eudesmol (11.57%), while *Piper vicosanum* contained α -eudesmol (17.35%) and α -copaene (12.34%). *Piper arboreum* var. *arboreum* and *Piper arboretum* var. *hirtelum* had a major *E*-caryophyllene (13.71% and 13.22%) in common, but a significant difference (>5%) in the relative percentage of bicyclogermacrene (18.50 % and 5.32%) and α -cadinol (2.40% and 8.89%), respectively. Studies such as the one presented in this manuscript serve as a means to prevent adulteration

and to describe the different chemophenetic phenomena of this species.

The majority of studies on EOs with a high content of *E*-nerolidol can be found in the Brazilian Atlantic Forest. For example, *P. aduncum* L. (80.6 – 82.5% content) [28, 29]; *P. clausenianum* (Miq.) C. DC. (81.4 – 83.3% content) [30]; *Piper diospyrifolium* Kunth (10.0% content) [27] and *P. gaudichaudianum* Kunth (22.1% - 22.4% content) [31-32, 14]. Chan and co-workers [33] state that *E*-nerolidol is widely used in the industry, including in the production of cosmetics, food, and pharmaceutical products. *Piper* species that are abundant in *E*-nerolidol have already been demonstrated to exhibit a variety of biological functions, such as antileishmanial (promastigotes of *Leishmania amazonensis*, IC₅₀ 30.24 μ g/ mL) and antifungal (*Candida albicans*, MIC 0.2 – 1.26%) for *P. clausenianum* [30, 34-35]; and cytotoxic (V79 Chinese hamster lung cells, IC₅₀ 4.0 μ g/ mL) for *P. gaudichaudianum* [31].

The compounds δ -cadinene, α -copaene, and valencene are described in the literature for their diverse biological activities when present in the essential oils of *Piper* [4-5, 7, 11, 14, 17]. These activities include antimicrobial, anti-inflammatory, antioxidant, and anticancer properties. Some studies also suggest

that these compounds may possess analgesic and sedative properties, in addition to their use in the fragrance and flavor industry due to their pleasant aroma [14, 17, 22-23].

4. Conclusions

The essential oil of the *P. robustipedunculum* leaf can be a source of vegetable raw material for future employability in the pharmaceutical, cosmetic, food and pesticide industries, since it is sesquiterpene-rich. Also, the chemical composition of this endemic species from the Atlantic Forest of Northeast Brazil, a biome high threatened by human action, can assist in studying to understand ecological, chemotaxonomics and chemophenetics issues considering Piperaceae species.

Authors' contributions

Conceptualization, G.A. Q. and Y. J. R., Methodology, Y. J. R., E. F. G. and D. L. M.; Formal Analysis, G.A. Q., D. B. M., C. C.-O. and J. R. S. F.; Investigation, Y. J. R., G.A. Q. and I.C.F.; Resources, Y. J. R., D. B. M., C. C.-O., J. R. S. F., G.A. Q. and I.C.F.; Writing – Original Draft Preparation, Y. J. R. and I.C.F.; Writing – Review & Editing, Y. J. R., I.C.F. and D.L.M.

Funding

This research was funded by CNPq (Conselho Nacional de Pesquisas e Desenvolvimento Científico e Tecnológico e Inovação) - Brazil (PROEP n. 407845/2017); FAPERJ (Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro) – Brazil, (APQ1 n. NE26/210.245/2019), (CNE n. 201.211/2022).

Conflicts of interest

The authors declare no conflict of interest. The funding entity had no role in the design of the study, in the collection, analysis, or interpretation of data, in the writing of the manuscript, or in the decision to publish the results.

References

1. Flora do Brasil. Jardim Botânico do Rio de Janeiro. Available in <<http://floradobrasil.jbrj.gov.br/>>. Acesso em 9 outubro 2022.
2. Wanke, S.; Samain, M.S.; Vanderschaeva, L.; Mathieu, G.; Goetghebeur, P.; Neinhuis, C. Phylogeny of the genus *Peperomia* (Piperaceae) inferred from the trnk/matk region (cpDNA). *Plant. Biol.* 2006, 8, 1, 93-102. <https://doi.org/10.1055/s-2005-873060>.
3. Jaramillo, M.A.; Callejas, R.; Davidson, C.; Smith, J.F.; Stevens, A.C.; Tepe, E.J. A phylogeny of the tropical genus *Piper* using ITS and the chloroplast intron psbJ–petA. *Syst. Bot.* 2008, 33, 4, 647-660. <https://doi.org/10.1600/036364408786500244>.
4. Yadav, V.; Krishnan, A.; Vohora, D. A systematic review on *Piper longum* L.: Bridging traditional knowledge and pharmacological evidence for future translational research. *J. Ethnopharmacol.* 2019, 247, 112255. <https://doi.org/10.1016/j.jep.2019.112255>.
5. Takooree, H.; Aumeeruddy, M.Z.; Rengasamy, K.R.; Venugopala, K.N.; Jeewon, R.; Zengin, G.; Mahomoodally, M.F. A systematic review on black pepper (*Piper nigrum* L.): from folk uses to pharmacological applications. *Crit. Rev. Food Sci. Nutr.* 2019, 59, 210-243. <https://doi.org/10.1080/10408398.2019.1565489>.
6. Pereira, L.A.; Lima, R.B.; Guimarães E.F.; Almeida, M.Z.; Monteiro, E.D.C.Q.; Sobrinho, F.D.A.P. Plantas medicinais de uma comunidade quilombola na Amazônia Oriental: Aspectos utilitários de espécies das famílias Piperaceae e Solanaceae. *Rev. Bras. Agroecol.* 2007, 2, 2, 1385-1388.
7. Salehi, B.; Zakaria, Z.A.; Gyawali, R.; Ibrahim, S.A.; Rajkovic, J.; Shinwari, Z.K.; Valussi, M. *Piper* species: A comprehensive review on their phytochemistry, biological activities and applications. *Molecules* 2019, 24, 7, 1364. <https://doi.org/10.3390/molecules24071364>.
8. Gogosz, A.M.; Boeger, M.R.T.; Negrelle, R.R.B.; Bergo, C. Anatomia foliar comparativa de nove espécies do gênero *Piper* (Piperaceae). *Rodriguésia*. 2012, 63, 2, 405-417. <https://doi.org/10.1590/S2175-78602012000200013>.
9. Silva, R.J.F.; Faial, K.D.C.F.; Mendonça, M.S.D. Pharmacognostical characterization of *Piper arboreum* var. *arboreum* and *P. tuberculatum* (Piperaceae). *Acta Amaz.* 2016, 46, 2, 195-208. <https://doi.org/10.1590/1809-4392201504422>.
10. Moreira, D.L.; Souza, P.O.; Kaplan, M.A.C.; Pereira, N.A.; Cardoso, G.L.; Guimarães, E.F. Effect of leaf essential oil from *Piper solmsianum* C. DC. in mice behaviour. *An. Acad. Bras. Ciênc.* 2001, 1, 73-78. <https://doi.org/10.1590/S0001-37652001000100004>.
11. Scott, I.M.; Jensen, H.R.; Philogène, B.J.; Arnason, J.T. A review of *Piper* spp. (Piperaceae) phytochemistry, insecticidal activity and mode of action. *Phytochem.* 2008, 7, 1, 65-75. <https://doi.org/10.1007/s11101-006-9058-5>.
12. Oliveira, G.L.; Cardoso, S.K.; Lara-Junior, C.R.; Vieira, T.M.; Guimarães, E.F.; Figueiredo, L.S.; Moreira, D.L.; Kaplan, M.A.C. Chemical study and larvicidal activity against *Aedes aegypti* of essential oil of *Piper aduncum* L.

- (Piperaceae). An. Acad. Bras. Ciênc. 2013, 85, 4, 1227-1234. <https://doi.org/10.21577/1984-6835.20190107>.
13. Ramos, Y.J.; Moreira, D.L. Seasonal Study of Essential Oil from Aerial Parts of *Peperomia galioides* Kunth (Piperaceae). Rev. Virtual Quim. 2019, 11, 5, 1540-1550. <https://doi.org/10.21577/1984-6835.20190107>.
 14. Ramos, Y.J.; Gouvêa-Silva, J.G.; Brito Machado, D.; Felisberto, J.S.; Pereira, R. C.; Sadgrove, N.J.; Moreira, D.L. Chemophenetic and Chemodiversity Approaches: New Insights on Modern Study of Plant Secondary Metabolite Diversity at Different Spatiotemporal and Organizational Scales. Rev. Bras. Farmacog. 2022, 1-24. <https://doi.org/10.1007/s43450-022-00327-w>.
 15. Guimarães, E.F.; Monteiro, D. Piperaceae of the Biological Reserve of Poço das Antas, Silva Jardim, Rio de Janeiro, Brazil. *Rodriguésia* 2006, 57, 3, 569-589. <https://doi.org/10.1590/2175-7860200657312>.
 16. Oliveira, R.A.; Assis, A.M.A.D.; Silva, L.A.M.; Andrioli, J.L.; Oliveira, F.F. Chemical profile and antimicrobial activity of essential oil of *Piper ilheusense*. Chem. Nat. Comp. 2016, 52, 2, 331-333. <https://doi.org/10.1007/s10600-016-1634-3>.
 17. Silva, J.K.; Trindade, R.; Alves, N.S.; Figueiredo, P.L.; Maia, J.G.S.; Setzer W.N. Essential oils from neotropical *Piper* species and their biological activities. Int. J. Mol. Sci. 2017, 18, 12, 2571. <https://doi.org/10.3390/ijms18122571>.
 18. Queiroz, G.A.; Guimarães, E.F.; Sakuragui, C.M. First record of *Piper robustipedunculum* Yunck. (Piperaceae) in the state of Pernambuco, Brazil. Check List. 2020, 16, 1149. <https://doi.org/10.15560/16.5.1149>.
 19. Dool, H.V.D.; Kratz, P.A. Generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. J. Chromatogr. 1963, 1, 463-471. [https://doi.org/10.1016/s0021-9673\(01\)80947-x](https://doi.org/10.1016/s0021-9673(01)80947-x).
 20. Ramos, Y.J.; Brito Machado, D.; Queiroz, G.A.; Guimarães, E.F.; Defaveri, A.C.A.; Moreira, D. L. Chemical composition of the essential oils of circadian rhythm and of different vegetative parts from *Piper mollicomum* Kunth-A medicinal plant from Brazil. Biochem. Syst. Ecol. 2020, 92, 104116. <https://doi.org/10.1016/j.bse.2020.104116>.
 21. Adams, R.P. Identification of essential oil components by gas chromatography/mass spectroscopy. Ed. IV; Allured Bus. Mul, Carol Stream: Illinois, USA, 2009.
 22. Filly, A.; Fernandez, X.; Minuti, M.; Visinoni, F.; Cravotto, G.; Chemat, F. Solvent-free microwave extraction of essential oil from aromatic herbs: from laboratory to pilot and industrial scale. Food Chem. 2014, 150 1, 193-198. <https://doi.org/10.1016/j.foodchem.2013.10.139>.
 23. Kemprai, P.; Bora, P.K.; Mahanta, B.P.; Sut, D.; Saikia, S.P.; Banik, D.; Haldar, S. *Piper betleoides* C. DC.: Edible source of betel-scented sesquiterpene-rich essential oil. Flavour Fragr. J. 2020, 35, 1, 70-78. <https://doi.org/10.1002/ffj.3537>.
 24. Santos, P.R.D.; Moreira, D.; Guimarães, E.F.; Kaplan, M.A.C. Essential oil analysis of 10 Piperaceae species from the Brazilian Atlantic forest. Phytochem. 2001, 58, 4, 547-551. [https://doi.org/10.1016/s0031-9422\(01\)00290-4](https://doi.org/10.1016/s0031-9422(01)00290-4).
 25. Miquel, F.A.G. *Systema Piperacearum*. Kramer, Rotterdam, 1843-1844.
 26. Yuncker, T.G. New species of Piperaceae from Brazil. *Bol. Inst. Bot.* 1966, 3, 1-370.
 27. Pereira, R.A.; Ramos Y.J.; Queiroz, G.A.; Guimarães E.F.; Defaveri, A.C.A.; Moreira, D.L. Chemodiversity of Essential Oils in *Piper* L.(Piperaceae) Species from the Restinga of Marambaia Island, Rio de Janeiro-RJ, Brazil. Rev. Virtual Quím. 2021, 13, 5, 1203-1215. <https://doi.org/10.21577/1984-6835.20210067>.
 28. Ceole, L.F.; Cardoso, M.D; Soares, M.J. Nerolidol, the main constituent of *Piper aduncum* essential oil, has anti-*Leishmania braziliensis* activity. Parasitol. 2017, 144, 9,1179-1190. <https://doi.org/10.1017/S0031182017000452>.
 29. Villamizar, L.H.; Cardoso, M.D.; de Andrade, J.; Teixeira, M.L.; Soares, M.J. Linalool, a *Piper aduncum* essential oil component, has selective activity against *Trypanosoma cruzi* trypomastigote forms at 4 °C. Mem. Inst. Oswaldo Cruz. 2017, 112, 131-139. <https://doi.org/10.1590/0074-02760160361>.
 30. Marques, A.M.; Barreto, A.L.S.; Batista, E.M.; Curvelo, J.A.R.; Velozo, L.S.M.; Moreira, D.L.; Guimarães, E.F.; Soares, R.M.A.; Kaplan, M.A.C. Chemistry and biological activity of essential oils from *Piper claussonianum* (Piperaceae). Nat. Prod. Commun. 2010, 5, 1837-1840. <https://doi.org/10.1177/1934578X1000501131>.
 31. Péres, V.F.; Moura, D.J.; Sperotto, A.R.M.; Damasceno, F.C.; Caramão, E.B.; Zini, C.A.; Saffi J. Chemical composition and cytotoxic, mutagenic and genotoxic activities of the essential oil from *Piper gaudichaudianum* Kunth leaves. Food Chem. Toxicol. 2009, 47, 2389-2395. <https://doi.org/10.1016/j.fct.2009.06.035>.
 32. Sperotto, A.R.M.; Moura, D.J.; Péres, V.F.; Damasceno, F.C.; Caramão, E.B.; Henriques, J.A.P.; Saffi, J. Cytotoxic mechanism of *Piper gaudichaudianum* Kunth essential oil and its major compound nerolidol. Food Chem. Toxicol. 2013, 57, 57-68. <https://doi.org/10.1016/j.fct.2013.03.013>.
 33. Chan, W.K.; Tan, L.T.H; Chan, K.G.; Lee, L.H.; Goh, B.H. Nerolidol: A Sesquiterpene Alcohol with Multi-Faceted Pharmacological and Biological Activities. Molecules. 2016, 21, 5, 529. <https://doi.org/10.3390/molecules>

- 21050529.
34. Marques, A.M.; Barreto, A.L.S.; Curvelo, J.A.D.R.; Romanos, M.T.V.; Soares, R.M.D.A.; Kaplan, M.A.C. Antileishmanial activity of nerolidol-rich essential oil from *Piper claussonianum*. Rev. Bras. Farmacogn. 2011, 21, 5, 908-914. <https://doi.org/10.1590/S0102-695X2011005000157>.
35. Curvelo, J.A.R.; Marques, A.M.; Barreto, A.L.S.; Romanos, M.T.V.; Portela, M.B.; Kaplan, M.A.C.; Soares, R.M.A.; A novel nerolidol-rich essential oil from *Piper claussonianum* modulates *Candida albicans* biofilm. J. Med. Microbiol. 2014. 63, 5, 697-702. <https://doi.org/10.1099/jmm.0.063834-0>.